

Acknowledgements

Genotyping for this analysis was funded by the Alaska Fisheries Science Center, National Marine Fisheries Service, NOAA. We are very grateful to Bill Templin of the ADFG Gene Conservation Laboratory for sharing the SNP baseline, providing suggestions and advice regarding the analysis, and reviewing this report. We are also grateful to the outstanding help from the North Pacific Groundfish Observer Program including Lisa Thompson, Craig Faunce, Jennifer Cahalan, Paul Wilkins, and the many participating observers who helped fulfill our request for genetic samples. MALDI-TOF genotyping and assay design was performed in collaboration with the genotyping core facility at the University of Arizona Genetics Core (Libby Serbi and Ryan Sprissler). Andrew Gray, Chris Kondzela, Tyler McCraney, Frank Thrower and Colby Marvin helped review the report. This document has been peer reviewed by internal and external reviewers.

References

- ADFG (Alaska Department of Fish and Game. (2003). SPAM Version 3.7: Statistics Program for Analyzing Mixtures. Alaska Department of Fish and Game, Commercial Fisheries Division, Gene Conservation Lab, Anchorage, AK.
- Gabriel, S., Ziaugra, L., and Tabbaa, D. (2009). SNP genotyping using the Sequenom MassARRAY iPLEX platform. *Current Protocols in Human Genetics Chapter 2, Unit 2 12.*
- Guyon, J.R., Guthrie, C.M., and Nguyen, H. (2010). Genetic Stock Composition Analysis of Chinook Salmon Bycatch Samples from the 2008 Bering Sea Pollock Fishery, Report to the North Pacific Fisheries Management Council. (Juneau, AK, National Marine Fisheries Service, Alaska Fisheries Science Center, Auke Bay Laboratories), pp. 32.
- Locke, G. (2010). Commerce Secretary Gary Locke announces "Fishery Failure" determination for Alaska Chinook salmon. In Commerce News release, January 15, 2010, U.S. Department of Commerce, Washington, DC.
- NMFS (National Marine Fisheries Service). (2009). Bering Sea Chinook salmon bycatch management - Volume 1, Final Environmental Impact Statement, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, Alaska Regional Office, Juneau, AK.
- NMFS (National Marine Fisheries Service). (2010). BSAI Chinook salmon mortality estimates, 1991-present, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, Alaska Regional Office, Juneau, AK.
http://www.fakr.noaa.gov/sustainablefisheries/inseason/chinook_salmon_mortality.pdf
- Peakall, R., and Smouse, P.E. (2006). Genalex 6: genetic analysis in Excel. Population genetic software for teaching and research. *Molecular Ecology Notes* 6, 288-295.
- Pella, J., and Geiger, H.J. (2009). Sampling considerations for estimating geographic origins of Chinook salmon bycatch in the Bering Sea pollock fishery. Alaska Department of Fish and Game Special Publication No. SP 09-08.
- Pella, J., and Masuda, M. (2001). Bayesian methods for analysis of stock mixtures from genetic characters. *Fishery Bulletin* 99, 151-167.